

Clinical Policy: Lutetium Lu 177 Dotatate (Lutathera)

Reference Number: CP.PHAR.384

Effective Date: 05.22.18

Last Review Date: 08.18

Line of Business: Medicaid, HIM-Medical Benefit

[Coding Implications](#)[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Lutetium Lu 177 dotatate (Lutathera[®]) is a radiolabeled somatostatin analog.

FDA Approved Indication(s)

Lutathera is indicated for the treatment of somatostatin receptor-positive gastroenteropancreatic neuroendocrine tumors (NETs), including foregut, midgut, and hindgut NETs in adults.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation[®] that Lutathera is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria**A. Neuroendocrine Tumors (must meet all):**

1. Diagnosis of a somatostatin receptor-positive NET of one of the following origins (a or b):
 - a. Gastrointestinal tract or pancreas;
 - b. Lung or thymus (off-label);
2. Prescribed by or in consultation with an oncologist;
3. Age \geq 18 years;
4. Disease is metastatic or locally advanced, and unresectable;
5. Member experienced disease progression while on a long-acting somatostatin analog (e.g., octreotide, lanreotide);
6. Member has not received \geq 4 doses of Lutathera;
7. Dose does not exceed 7.4 GBq (200 mCi) every 8 weeks, up to a total of 4 doses.

Approval duration: 32 weeks (no more than 4 total doses)

B. Pheochromocytoma/Paraganglioma (off-label) (must meet all):

1. Diagnosis of a somatostatin receptor-positive pheochromocytoma/paraganglioma;
2. Prescribed by or in consultation with an oncologist;
3. Disease is metastatic or locally advanced, and unresectable;
4. Member has not received \geq 4 doses of Lutathera;
5. Dose does not exceed 7.4 GBq (200 mCi) every 8 weeks, up to a total of 4 doses.

Approval duration: 32 weeks (no more than 4 total doses)

C. Other diagnoses/indications

1. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): HIM.PHAR.21 for health insurance marketplace and CP.PMN.53 for Medicaid.

II. Continued Therapy

A. All Indications in Section I (must meet all):

1. Currently receiving medication via Centene benefit, or documentation supports that member is currently receiving Lutathera for a covered indication;
2. Member is responding positively to therapy;
3. Member has not received ≥ 4 doses of Lutathera;
4. If request is for a dose increase, new dose does not exceed 7.4 GBq (200 mCi) every 8 weeks, up to a total of 4 doses.

Approval duration: 32 weeks (no more than 4 total doses)

B. Other diagnoses/indications (must meet 1 or 2):

1. Currently receiving medication via Centene benefit and documentation supports positive response to therapy.
Approval duration: Duration of request or 6 months; or
2. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): HIM.PHAR.21 for health insurance marketplace and CP.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – HIM.PHAR.21 for health insurance marketplace and CP.PMN.53 for Medicaid or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

CT: computed tomography	NCCN: National Comprehensive Cancer Network
FDA: Food and Drug Administration	PET: positron emission tomography
NET: neuroendocrine tumor	

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
Somatuline [®] Depot (lanreotide)	120 mg SC every 4 weeks	120 mg/month

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
Sandostatin [®] LAR Depot (octreotide LAR)*	30 mg IM once monthly (20 mg may be used for pancreatic NETs)	30 mg/month

Therapeutic alternatives are listed as Brand name[®] (generic) when the drug is available by brand name only and generic (Brand name[®]) when the drug is available by both brand and generic.

**Off-label for the treatment of NETs (octreotide is only FDA-approved for the treatment of symptoms associated with carcinoid tumors) – NET dosing recommendations are per the NCCN guidelines*

Appendix C: Contraindications

Not applicable

Appendix D: General Information

- Somatostatin receptor expression can be detected by somatostatin receptor-based imaging, which includes ⁶⁸Ga-dotatate PET/CT (preferred per the NCCN) and somatostatin receptor scintigraphy.
- The NCCN Neuroendocrine and Adrenal Tumors guidelines recommend the use of Lutathera:
 - For somatostatin receptor-positive bronchopulmonary/thymus, gastrointestinal, and pancreatic NETs that have progressed following therapy with octreotide or lanreotide and are locoregionally advanced or have distant metastases (category 2A, except for mid-gut tumors [category 1]); and
 - For the primary treatment of somatostatin receptor-positive pheochromocytoma/ paraganglioma that is locally unresectable or has distant metastases (category 2A).
- Use of Lutathera with long-acting somatostatin analogs:
 - Before initiating Lutathera: Long-acting somatostatin analogs (e.g., long-acting octreotide) should be discontinued for at least 4 weeks prior to initiation of Lutathera. Short-acting octreotide can be administered as needed up to 24 hours prior to initiating Lutathera.
 - During Lutathera treatment: Long-acting octreotide 30 mg should be administered intramuscularly between 4 to 24 hours after each Lutathera dose. Long-acting octreotide should not be administered within 4 weeks of each subsequent Lutathera dose. Short-acting octreotide may be given for symptomatic management during Lutathera treatment, but must be withheld for at least 24 hours before each Lutathera dose.
 - Following Lutathera treatment: Long-acting octreotide 30 mg intramuscularly should be continued every 4 weeks after completing Lutathera until disease progression or for up to 18 months following treatment initiation.

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
GEP-NET	7.4 GBq (200 mCi) IV every 8 weeks for a total of 4 doses	See regimen
NET of lung or thymus origin, pheochromocytoma, paraganglioma*		

**Off-label – dosing recommendations are per the NCCN guidelines*

VI. Product Availability

Single-dose vial for injection: 370 MBq/mL (10 mCi/mL)

VII. References

1. Lutathera Prescribing Information. Millburn, NJ: Advanced Accelerator Applications USA, Inc.; January 2018. Available at: <https://www.lutathera.com>. Accessed May 7, 2018.
2. National Comprehensive Cancer Network. Neuroendocrine and Adrenal Tumors. Version 2.2018. Available at: https://www.nccn.org/professionals/physician_gls/pdf/neuroendocrine.pdf. Accessed May 7, 2018.
3. National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at: http://www.nccn.org/professionals/drug_compendium. Accessed May 7, 2018.
4. Strosberg J, El-Haddad G, Wolin E, et al. Phase 3 trial of ¹⁷⁷Lu-dotatate for midgut neuroendocrine tumors. N Engl J Med. 2017; 376(2): 125-135.
5. Brabander T, van der Zwan WA, Teunissen JJM, et al. Long-term efficacy, survival, and safety of [¹⁷⁷Lu-DOTA⁰,Tyr³]octreotate in patients with gastroenteropancreatic and bronchial neuroendocrine tumors. Clin Cancer Res. 2017; 1-8.

Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
C9399	Unclassified drugs or biologicals
A9699	Radiopharmaceutical, therapeutic, not otherwise classified

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy created.	05.22.18	08.18

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note:

For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

For Health Insurance Marketplace members, when applicable, this policy applies only when the prescribed agent is on your health plan approved formulary. Request for non-formulary drugs must be reviewed using the formulary exception policy.

©2018 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed,

displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene[®] and Centene Corporation[®] are registered trademarks exclusively owned by Centene Corporation.